

What is claimed is:

1. A stable, sterile, and injectable aqueous dispersion of a water-insoluble microdroplet matrix of mean diameter from about 50 nm to about 1000 nm consisting essentially of
 - (a) between about 1% to about 15% of propofol;
 - (b) between about 1% to about 8% of a propofol soluble diluent;
 - (c) between about 0.5% to about 5% of a surface stabilizing amphiphilic agent;
 - (d) of a pharmaceutically acceptable water-soluble polyhydroxy additive that acts as a tonicity modifier; and
 - (e) provided the ratio of propofol to diluent is about 1:4 to about 1:0.1 and the ratio of propofol to amphiphilic agent is about 1:0.8 to about 1:2.5, and the composition has a viscosity of from about 0.8 to about 15 centipoise, wherein the formulation
 - prevents microbial growth, defined as no more than 0.5 log increase from the initial inoculum, of each of *Staphylococcus aureus* (ATCC 6538), *Escherichia coli* (ATCC 8739 and ATCC 8454), *Pseudomonas aeruginosa* (ATCC 9027), *Candida albicans* (ATCC 10231), and *Aspergillus niger* (ATCC 16403) for at least 7 days as measured by a test wherein a washed suspension of each said organism is added to a separate aliquot of a formulation at approximately 1000 colony forming units (cfu) per mL, at a temperature in the range 20-25°C, whereafter said aliquots are incubated at 20-25°C and are tested for viability of the microorganisms in the inoculated formulation as determined by counting the colonies of

said organism after 24, 48 hours and 7 days or other suitable length of time;

- results in little or no irritation at the site of injection as evidenced by a test wherein said composition is administered as a single daily bolus injection of 12.5 mg/kg, given on the basis of body weight, for 2 or 3 successive days over a period of approximately 30 seconds, in the caudal vein of a rat such that no visual increase in the diameter of the rat tail is noted after 48hrs post injection.

2. The composition of claim 1 wherein the surface stabilizing amphiphilic agent is one or more natural or synthetic surface modifiers selected from ionizable or non-ionizable phospholipids or cholesterol or a mixture of these amphiphilic agents, the total quantity of the amphiphilic agents being such that the ratio of propofol to amphiphilic agent is about 1:0.8 to about 1:2.5 and the types and quantities of the individual amphiphilic agents are selected to provide that:
 - (i) the composition has a non-existent or minimum potential for hemolysis of human or animal blood, and
 - (ii) the irritation to the tissues at the site of injection is either non-existent or minimized, and
 - (iii) the composition elicits an anesthetic effect in warm-blooded animal and human subjects upon intravenous administration.
3. The composition of claim 1 wherein the propofol-soluble diluent is a synthetic or natural fatty acid, triglyceride thereof or other suitable ester or a mixture thereof.
4. The composition of claim 1 wherein the ratio of propofol to the amount of propofol-soluble diluent is from about 1:3 to about 1:0.5.

5. The composition of claim 1 wherein the ratio of propofol to the amount of propofol-soluble diluent is from about 1:2 to about 1:1.
6. The composition of claim 1 wherein the propofol-soluble diluent is a mixture of medium-chain triglyceride and vegetable oil.
7. The composition of claim 6 wherein the ratio of medium-chain triglyceride to vegetable oil is from 1:3 to 3:1.
8. The composition of claim 1 wherein the water-insoluble matrix consists of a mixture of the amphiphilic agents of claim 3 and propofol-soluble diluents of claims 4-7 and propofol.
9. The composition of claim 1 wherein the composition contains about 2% to about 10% of propofol.
10. The composition of claim 1 wherein the composition contains a pharmaceutically acceptable water-soluble polyhydroxy additive that provides the propofol containing dispersion with an osmolality of about 250 to about 700 milliosmolal.
11. The composition of claim 1 wherein the osmolality is about 300 to about 500 milliosmolal.
12. The composition of claim 1 wherein the viscosity is from about 2 to about 5 centipoise.

13. The method of reducing or substantially completely eliminating irritation upon injection of formulations containing propofol by administering a stable, sterile, and antimicrobial, aqueous dispersion of water-insoluble microdroplet matrix of mean diameter from about 50 nm to about 1000 nm consisting essentially of about 1% to about 15% of propofol as the active ingredient, up to about 7% of a propofol soluble diluent, and about 0.8% to about 4% of a surface stabilizing amphiphilic agent, and the aqueous phase of the composition consisting of a pharmaceutically acceptable water-soluble polyhydroxy tonicity modifier, the composition being devoid of additional bactericidal or bacteriostatic preservative agents, provided the ratio of propofol to diluent is about 1:4 to about 1:0.1 and the ratio of propofol to amphiphilic agent is about 1:0.8 to about 1:2.5, and the composition has a viscosity of from about 0.8 to about 15 centipoise.
14. The method of inducing anesthesia comprising administering to a subject in need of same an anesthetic inducing amount of a stable, sterile, and antimicrobial injectable aqueous dispersion of a water-insoluble microdroplet matrix of mean diameter from about 50 nm to about 1000 nm consisting essentially of about 1% to about 15% of propofol as the active ingredient, up to about 7% of a propofol soluble diluent, and about 0.8% to about 4% of a surface stabilizing amphiphilic agent, and the aqueous phase of the composition consisting of a pharmaceutically acceptable water-soluble polyhydroxy tonicity modifier, the composition being devoid of additional bactericidal or bacteriostatic preservative agents, provided the ratio of propofol to diluent is about 1:4 to about 1:0.1 and the ratio of propofol to amphiphilic agent is about 1:0.8 to about 1:2.5, and the composition has a viscosity of from about 0.8 to about 15 centipoise.

REFERENCES

- Ardulno, M. J., Bland, L. A., McAllister, S. K., Agüero, S. M., Villarino, M. E., McNeil, M. M., Jarvis, W. R. and Favero, M. S. (1991) "Microbial Growth and Endotoxin Production in the Intravenous Anesthetic Propofol" *Inf. Control Hosp. Epidem.*, 12(9), 535-539.
- J. Babl, A. Doenicke, V. Monch (1995) "New propofol LCT/MCT fat emulsions as solvent. Approach to reducing pain on injection of propofol." *Eur Hosp Pharmacy*, 1: 15-21.
- Cairns *et al.* (1996) "Tolerance of mixed lipid emulsion in neonates: effect of concentration." *Arch Dis Child Fetal Neonatal Ed* 75(2): F113-6.
- Crowther, J., Hrazdil, J., Jolly, D. T., Galbraith, J. C., Greacen, M. and Grace, M. (1996) "Growth of Microorganisms in Propofol, Thiopental and a 1:1 Mixture of Propofol and Thiopental" *Anesth. and Anal.* 82, 475-478.
- A.W. Doenicke, J. Babl, W. Kellermann, J. Rau, M.F. Roizen (1996) "Reducing pain during propofol injection: the role of the solvent." *Anesth Analg* 82: 472-4.
- A.W. Doenicke, J. Babl, U. Klotz, J. Kugler, M. O'Connor, J. Rau, M.F. Roizen (1997) "Pharmacokinetics and pharmacodynamics of propofol in a new solvent." *Anesth Analg* 85: 1399-403.
- A.F. Ghouri, M.A. Ramirez Ruiz, and P.F. White (1994) "Effect of flumazenil on recovery after midazolam and propofol sedation." *Anesthesiology* 81: 333-339.
- J.B. Glen and R. James "2,6-Diisopropylphenol as an anaesthetic agent." US Patent 4,056,635. Nov 1, 1977
- J.B. Glen and R. James "Anaesthetic compositions containing 2,6-diisopropylphenol." US Patent 4,452,817. Jun 5, 1984
- J.B. Glen and R. James "Pharmaceutical compositions." US Patent 4,798,846. Jan 17, 1989
- D.H. Haynes "Propofol microdroplet formulations." US Patent 5,637,625. Jun. 10, 1997.
- C. B. Jones and J.H. Platt "Propofol composition containing edetate." US Patent 5,714,520. Feb 3, 1998.
- C. B. Jones and J.H. Platt "Pharmaceutical compositions of propofol and edetate." US Patent 5,731,355. Mar 24, 1998.

H. G. Weder "Propofol Nanodispersions." PCT Patent Application WO9710814. March 27 1997.

[illegible]